antibodies -online.com





Datasheet for ABIN1386591

anti-LRRTM2 antibody (AA 81-200)



Overview

Quantity:	100 μL
Target:	LRRTM2
Binding Specificity:	AA 81-200
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LRRTM2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human LRRTM2
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target: LRRTM2

Target Details

J	
Alternative Name:	LRRTM2 (LRRTM2 Products)
Background:	Synonyms: KIAA0416, Leucine rich repeat transmembrane neuronal 2, Leucine rich repeat
	transmembrane neuronal 2 protein, Leucine rich repeat transmembrane protein 2, Leucine-rich
	repeat neuronal 2 protein, Leucine-rich repeat transmembrane neuronal protein 2, LRRN2,
	LRRT2_HUMAN, Lrrtm2.
	Background: The leucine-rich (LRR) repeat is a 20-30 amino acid motif that forms a
	hydrophobic Alpha/Beta horseshoe fold, allowing it to accommodate several leucine residues
	within a tightly packed core. All LRR repeats contain a variable segment and a highly conserved
	segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. The LRRTM
	protein family plays a role in the regulation of various cellular events during nervous system
	development. Localizing predominantly to the nervous system, LRRTM family members are
	known to exhibit synaptogenic activity. LRRTM2 (leucine rich repeat transmembrane neuronal
	2), also known as LRRN2, is a 516 amino acid single-pass type I membrane protein involved in
	the development maintenance of the vertebrate nervous system. Expressed in kidney and
	neuronal tissues, LRRTM2 contains ten LRR repeats and belongs to the LRRTM family.
	LRRTM2 is encoded by a gene that maps to human chromosome 5q31.2.
Pathways:	Synaptic Membrane
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months