

Datasheet for ABIN1386623
anti-POU3F3 antibody (AA 351-450)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	POU3F3
Binding Specificity:	AA 351-450
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This POU3F3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Brain1/POU3F3
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	POU3F3
Alternative Name:	Brain1/POU3F3 (POU3F3 Products)

Target Details

Background: Synonyms: Brain 1, Brain specic homeobox/POU domain protein 1, Brain-1, Brain-specific homeobox/POU domain protein 1, Brain1, BRN 1, Brn-1, BRN1, class 3, Oct-8, Octamer-binding protein 8, Octamer-binding transcription factor 8, OTF 8, OTF-8, OTF8, PO3 F3, PO3F 3, PO3F3, PO3F3_HUMAN, POU class 3 homeobox 3, POU domain, POU domain, class 3, transcription factor 3, POU3 F3, POU3F 3, Pou3f3, RHS 1, Rhs 2, RHS1, Rhs2, Skin 1, Skin1, transcription factor 3.

Background: The Brn family of transcription factors are found in a highly restricted subset of neurons and are critical to the early embryonic development of the central nervous system. Brn-1 and Brn-2 are class III POU domain proteins. Expressed during the development of the forebrain and coexpressed in most layer II-V cortical neurons, Brn-1 and Brn-2 appear to critically control the initiation of radial migration of cortical neurons. Brn-2 is thought to be involved in smooth muscle cell development and differentiation. Brn-3 is a class IV POU domain protein. Three Brn-3 proteins have been described and are designated Brn-3a, Brn-3b and Brn-3c. Brn-3a has two functional transactivating domains, one at the amino terminus and one at the carboxy terminus. While Brn-3a and Brn-3c stimulate transcription, Brn-3b generally functions as a transcriptional repressor. However, Brn-3b, but not Brn-3a, has been shown to regulate the expression of the acetylcholine receptor.

Application Details

Application Notes: WB 1:300-5000
ELISA 1:500-1000
IHC-P 1:200-400
IHC-F 1:100-500
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
ICC 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months