antibodies -online.com





anti-PATJ antibody (AA 1001-1200)



Overview

Quantity:	100 μL
Target:	PATJ (INADL)
Binding Specificity:	AA 1001-1200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PATJ antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PATJ
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	PATJ (INADL)
Alternative Name:	PATJ (INADL Products)

Target Details

Background:

Synonyms: Channel interacting PDZ domain protein, Cipp, FLJ26982, hINADL, Inactivation no after potential D like protein, INAD like, InaD like Drosophila, INAD like protein, INADL, INADL protein, PALS 1 associated tight junction protein, PALS1 associated tight junction protein, PATJ, PDZ domain protein, PDZ domain protein Drosophila inaD like, Post synaptic density 95 / discs large / zonula occludens 1 domain protein, Protein associated to tight junctions, INADL_HUMAN.

Background: The membranes of myelinating Schwann cells are joined by tight, gap and adherens junctions, all of which are found in regions of noncompact myelin: the paranodal loops, incisures of Schmidt-Lanterman and mesaxons. Tight junctions help establish polarity in mammalian epithelia by forming a physical barrier that separates apical and basolateral membranes. Pals-associated tight junction protein (PATJ), the human homolog of Drosophila Discs Lost, is differentially localized in myelinating Schwann cells. PATJ associates with Claudin-1, CRB1 (a transmembrane protein that plays a role in epithelial cell polarity and photoreceptor development), and Pals1 (a Lin-7 associated protein). The PATJ/Pals1/CRB1 complex can form a tripartite tight junction in epithelial cells crucial to their integrity.

Application Details

Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months