



[Go to Product page](#)

Datasheet for ABIN1386736
anti-KCNJ1 antibody (pSer44)

Overview

Quantity:	100 µL
Target:	KCNJ1
Binding Specificity:	pSer44
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KCNJ1 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human KCNJ1 around the phosphorylation site of Ser44
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Sheep,Horse
Purification:	Purified by Protein A.

Target Details

Target:	KCNJ1
Alternative Name:	KCNJ1 (KCNJ1 Products)

Target Details

Background: Synonyms: KCNJ1 phospho S44, p-KCNJ1 phospho S44, KCNJ1 phospho Ser44, p-KCNJ1 Ser44, p-ROM-Kphospho S44, KCNJ1 phospho S25, p-KCNJ1 phospho S25, KCNJ1 phospho Ser25, p-KCNJ1 Ser25, p-ROM-Kphospho S25, ROM K, ROM-K, inwardly rectifying subfamily J member 1, ATP regulated potassium channel ROM K, ATP sensitive inward rectifier potassium channel 1, ATP-regulated potassium channel ROM-K, ATP-sensitive inward rectifier potassium channel 1, Inward rectifier K⁺ channel Kir1.1, inwardly rectifying K⁺ channel, IRK1_HUMAN, KCNJ 1, KCNJ, Kcnj1, Kir 1.1, Kir1.1, Potassium channel, Potassium channel inwardly rectifying subfamily J member 1, potassium inwardly-rectifying channel J1, ROMK 1, ROMK 2, ROMK, ROMK1, ROMK2.

Background: Potassium channels are present in most mammalian cells, where they participate in a wide range of physiologic responses. The protein encoded by this gene is an integral membrane protein and inward-rectifier type potassium channel. It is activated by internal ATP and probably plays an important role in potassium homeostasis. The encoded protein has a greater tendency to allow potassium to flow into a cell rather than out of a cell. Mutations in this gene have been associated with antenatal Bartter syndrome, which is characterized by salt wasting, hypokalemic alkalosis, hypercalciuria, and low blood pressure. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008].

Application Details

Application Notes: ELISA 1:500-1000
IHC-P 1:200-400
IHC-F 1:100-500
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
ICC 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months