

Datasheet for ABIN1386746

anti-ZNF148 antibody (AA 303-380)



Go to Product page

_				
	۱۱ / ۱	rv		۱۸/
	' V '	 ı v	Ι.	v v

Quantity:	100 μL
Target:	ZNF148
Binding Specificity:	AA 303-380
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF148 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ZNF148/ZBP-89
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ZNF148
Alternative Name:	ZNF148/ZBP-89 (ZNF148 Products)

Target Details

Background:

Synonyms: Transcription factor ZBP-89, Transcription factor ZBP89, ZBP89, Zinc finger DNA binding protein 89, Zinc finger DNA-binding protein 89, Zinc finger protein 148, ZN148_HUMAN, ZNF 148, ZNF148.

Background: ZNF148 is a zinc finger transcription factor that is universally expressed. ZBP-89, a Kruppel-like repressor protein, is the silencer element binding factor for Vimentin. ZBP-89 has been shown to bind to GC-rich DNA elements in promoters for gastrin, ornithine decarboxylase and the cyclin-dependent kinase inhibitor p21 (also designated Cip1 or WAF1). ZBP-89 expression is induced by trans-retinoic acid or butyrate, which also induces terminal differentiation of colon cancer cells. ZBP-89 cooperates with histone acetyltransferase coactivator p300 in the regulation of p21, a cyclin-dependent kinase inhibitor whose associated gene is a target gene of p53. ZBP-89 also regulates cell proliferation, in part, through its ability to directly bind the p53 protein and retard its nuclear export. Elevated levels of ZBP-89 induce growth arrest and apoptosis in human gastrointestinal cells.

Gene ID:

7707

Application Details

Application Notes:	WB 1:300-5000

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

ICC 1:100-500

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months