antibodies - online.com







anti-RDH13 antibody (AA 101-200)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	RDH13
Binding Specificity:	AA 101-200
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RDH13 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RDH13
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	RDH13	

Target Details

Alternative Name:	RDH13 (RDH13 Products)	
Background:	Synonyms: RDH13, RDH13_HUMAN, Retinol dehydrogenase 13 all trans and 9 cis, Retinol	
	dehydrogenase 13 all trans/9 cis, Retinol dehydrogenase 13.	
	Background: RDH13, also known as all-trans and 9-cis retinol dehydrogenase 13 or SDR7C3, is	
	a 331 amino acid mitochondrial protein belonging to the short-chain	
	dehydrogenases/reductases (SDR) family. Widely expressed, mostly in eye, pancreas, placenta	
	and lung, RDH13 localizes on the outer side of the inner mitochondrial membrane. Related to	
	microsomal retinoid oxidoreductase RDH11, RDH13 is considered to be a major enzyme	
	among the RDH family of proteins. Catalytically active, RDH13 recognizes retinoids as	
	substrates and may function in retinoic acid production. RDH13 may function to protect the	
	mitochondria against oxidative stress. Leber congenital amaurosis (LCA) type 3, an inherited	
	autosomal recessive retinal disease, has been associated with defects of RDH13. LCA	
	represents the most common genetic cause of congenital visual impairment in infants and	
	children.	
Gene ID:	112724	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	

Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months