antibodies -online.com





Datasheet for ABIN1386823

anti-FBXW8 antibody (AA 181-280)



Overview

Quantity:	100 μL
Target:	FBXW8
Binding Specificity:	AA 181-280
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FBXW8 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffinembedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FBXW8/FBXO29
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

|--|

Target Details

Alternative Name:	FBXW8 (FBXW8 Products)
Background:	Synonyms: F box/WD repeat containing protein 8; F box and WD40 domain containing protein
	8; F-box only protein 29; F box only protein 29; FBW6; FBW8; FBX29; FBXO29; FBXW6;
	FBXW8_HUMAN.
	Background: FBXO29, also designated F-box/WD repeat-containing protein 8 (FBXW8), is a 598
	amino acid protein that contains one 40 amino acid F-box region, making it a member of the F-
	box family. FBXO29 also contains five WD repeats. F-box proteins are critical components of
	the SCF (Skp1-CUL-1-F-box protein) type E3 ubiquitin ligase complex and are involved in
	substrate recognition and recruitment for ubiquitination. F-box proteins are members of a large
	family that regulates cell cycle, immune response, signaling cascades and developmental
	programs by targeting proteins, such as cyclins, cyclin-dependent kinase inhibitors, IkB-å and -
	catenin, for degradation by the proteasome after ubiquitination. Functioning as a component of
	the SCF complex, FBXO29 is thought to recognize and bind to select phosphorylated proteins,
	thereby promoting their ubiquitination and subsequent degradation. FBXO29 exists as two
	isoforms as a result of alternative splicing events.
Gene ID:	26259
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months