



[Go to Product page](#)

Datasheet for ABIN1386842  
**anti-FNDC1 antibody (AA 21-120)**

### Overview

Quantity:	100 µL
Target:	FNDC1
Binding Specificity:	AA 21-120
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FNDC1 antibody is un-conjugated
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FNDC1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	FNDC1
Alternative Name:	FNDC1 ( <a href="#">FNDC1 Products</a> )

## Target Details

---

**Background:** Synonyms: Activation-associated cDNA protein, AGS8, Expressed in synovial lining protein, fibronectin type III domain containing 2, Fibronectin type III domain-containing protein 1, FNDC1, FNDC1\_HUMAN, FNDC2, MEL4B3.

Background: FNDC1, also known as MEL4B3, Ags8, Expressed in synovial lining protein and Activation-associated cDNA protein, is a 1888 amino acid secreted protein that contains five fibronectin type-III domains. FNDC1 is moderately expressed in skeletal muscle, pancreas, heart, kidney, spinal cord, ovary and lung. Expression of FNDC1 is induced in response to hypoxia in ventricular cardiomyocytes. Since FNDC1 interacts with G and G $\alpha$ , it is likely that FNDC1 is an activator for G-protein signaling. Though normally absent in healthy skin, FNDC1 expression is induced by TGF-beta signaling in skin tumors and psoriasis. There are two isoforms of FNDC1 that are produced as a result of alternative splicing events.

---

**Gene ID:** 84624

## Application Details

---

**Application Notes:** ELISA 1:500-1000  
IHC-P 1:200-400  
IHC-F 1:100-500  
IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

---

**Restrictions:** For Research Use only

## Handling

---

**Format:** Liquid

---

**Concentration:** 1  $\mu$ g/ $\mu$ L

---

**Buffer:** 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

---

**Preservative:** ProClin

---

**Precaution of Use:** This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

---

**Storage:** 4 °C,-20 °C

---

**Storage Comment:** Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

---

**Expiry Date:** 12 months