antibodies -online.com





anti-FBXL12 antibody (AA 171-270)



Overview

Quantity:	100 μL
Target:	FBXL12
Binding Specificity:	AA 171-270
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FBXL12 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FbxL12
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.
T 10.1.1	

Target Details

Target:	FBXL12

Target Details

Alternative Name:	FbxL12 (FBXL12 Products)
Background:	Synonyms: F box and leucine rich repeat protein 12, F box protein FBL12, F box/LRR repeat
	protein 12, FBXL 12, FLJ20188, FXL12_HUMAN.
	Background: F-box proteins are critical components of the SCF (Skp1-CUL-1-F-box protein) type
	E3 ubiquitin ligase complex and are involved in substrate recognition and recruitment for
	ubiquitination. They are members of a larger family of proteins that are involved in the
	regulation of a wide variety of cellular processes (including the cell cycle, immune responses,
	signaling cascades and developmental events) through the targeting of proteins, such as
	cyclins, cyclin-dependent kinase inhibitors, I°,B-å, and -catenin, for proteasomal degradation.
	FBL12 (F-box and leucine-rich repeat protein 12), also designated FBXL12, is a 326 amino acid
	protein that contains one F-box domain, five LRR (leucine-rich) repeats and exists as two
	alternatively spliced isoforms. FBL12 interacts with both Skp1 and CUL-1, and is encoded by a
	gene that maps to human chromosome 19p13.2.
Gene ID:	54850
Pathways:	SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
	ProClin
Preservative:	11001111

Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months