# antibodies -online.com





Go to Product page

# Datasheet for ABIN1386891

# anti-IQCK antibody

#### Overview

Quantity:	100 μL
Target:	IQCK
Reactivity:	Human
Neactivity.	Turnari
Host:	Rabbit
Clonality:	Polyclonal
Conjugato	This IOOK antihady is un conjugated
Conjugate:	This IQCK antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)),
	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
	inimationistochemistry (i dramin embedded occitions) (into (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human IQCK
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by Protein A.

## **Target Details**

Target:	IQCK
Alternative Name:	IQCK (IQCK Products)
Background:	Synonyms: FLJ20115, FLJ36575, IQ domain containing protein K, MGC35048, IQCK_HUMAN.  Background: Chromosome 16 encodes over 900 genes in approximately 90 million base pairs, makes up nearly 3 % of human cellular DNA and is associated with a variety of genetic

disorders. The GAN gene is located on chromosome 16 and, with mutation, may lead to giant axonal neuropathy, a nervous system disorder characterized by increasing malfunction with growth. The rare disorder Rubinstein-Taybi syndrome is also associated with chromosome 16, though through the CREBBP gene which encodes a critical CREB binding protein. Signs of Rubinstein-Taybi include mental retardation and predisposition to tumor growth and white blood cell neoplasias. Crohn's disease is a gastrointestinal inflammatory condition associated with chromosome 16 through the NOD2 gene. An association with systemic lupus erythematosis and a number of other autoimmune disorders with the pericentromeric region of chromosome 16 has led to the identification of SLC5A11 as a potential autoimmune modifier. alternatively splicing events.

Gene ID:

124152

## **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IF(IHC-P) 1:50-200

Restrictions: For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months