

[Go to Product page](#)

## Datasheet for ABIN1386896 **anti-PDZD2 antibody**

### Overview

Quantity:	100 µL
Target:	PDZD2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PDZD2 antibody is un-conjugated
Application:	Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PDZK3
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

### Target Details

Target:	PDZD2
Alternative Name:	PDZK3 ( <a href="#">PDZD2 Products</a> )
Background:	Synonyms: Activated in prostate cancer protein, AIPC, KIAA0300, Likely ortholog of rat plakophilin related armadillo repeat protein interacting PDZ protein antibody PAPIN, PDZ domain containing 2, PDZ domain containing 3, PDZ domain containing protein 2, PDZ domain

## Target Details

containing protein 3, PDZD2, PDZK 2, PDZK2, PDZK-2, PDZK 3, PDZK3, PDZK-3, PIN1, Plakophilin related armadillo repeat protein interacting PDZ protein like, Rat plakophilin related armadillo repeat protein interacting PDZ protein like, PDZD2\_HUMAN.

Background: Proteins containing PDZ domains have been shown frequently to bind the C-termini of transmembrane receptors or ion channels. They have also been shown to bind to other PDZ domain proteins and could possibly be involved in intracellular signalling. PDZK3 contains six PDZ domains and shares sequence similarity with pro-interleukin-16 (pro-IL-16). Like pro-IL-16, the encoded protein localizes to the endoplasmic reticulum and is thought to be cleaved by a caspase to produce a secreted peptide containing two PDZ domains. In addition, the PDZK3 gene is upregulated in primary prostate tumors and may be involved in the early stages of prostate tumorigenesis.

Gene ID: 23037

## Application Details

Application Notes: IHC-P 1:200-400  
IF(IHC-P) 1:50-200

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months