antibodies .- online.com







anti-RNF125 antibody



Overview

| Quantity: | 100 μL |
|--------------|--|
| Target: | RNF125 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This RNF125 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human TRAC1/RNF125 |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Purified by Protein A. |

Target Details

| Target: | RNF125 |
|-------------------|--|
| Alternative Name: | RNF125 (RNF125 Products) |
| Background: | Synonyms: T cell RING activation protein 1, E3 ubiquitin protein ligase RNF125, Ring finger protein 125, RNF 125, T cell ring protein identied in activation screen, TRAC 1, TRAC1, RN125_HUMAN. |

Target Details

Background: The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain this conserved domain are generally involved in the ubiquitination pathway of protein degradation. TRAC-1 (T-cell RING activation protein 1), also known as RNF125 (ring finger protein 125) or E3 ubiquitin-protein ligase RNF125, is a 232 amino acid novel E3 ubiquitin ligase that functions as a positive regulator in the T-cell receptor signaling pathway. Expressed predominantly in lymphoid tissues such as spleen, thymus and bone marrow, TRAC-1 has been found to inhibit pathogen-induced cytokine production and downregulates HIV replication.

Gene ID:

9611

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IF(IHC-P) 1:50-200

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| Expiry Date: | 12 months |