



[Go to Product page](#)

Datasheet for ABIN1387074  
**anti-MTA2 antibody**

### Overview

Quantity:	100 µL
Target:	MTA2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MTA2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MTA2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

### Target Details

Target:	MTA2
Alternative Name:	MTA2 ( <a href="#">MTA2 Products</a> )
Background:	Synonyms: Mata111, Metastasis associated 1 family member 2, Metastasis associated 1 like 1, Metastasis associated gene 1 like 1, Metastasis associated gene family member 2, Metastasis associated protein 2, Metastasis associated protein MTA 2, Metastasis associated protein

## Target Details

---

MTA2, Mmta2, MTA1 L1 protein, MTA1-L1 protein, MTA1L1, MTA1L1: metastasis-associated 1-like 1, P53 target protein in deacetylase complex, PID, MTA2\_HUMAN.

Background: MTA1 is a component of the NURD (nucleosome remodeling and histone deacetylation) complex, which is associated with ATP-dependent chromatin-remodeling and histone deacetylase activity. MTA1 functions in conjunction with other components of NURD to mediate transcriptional repression as it facilitates the association of repressor molecules with the chromatin. Structurally, MTA1 contains a single SH3-binding motif and a zinc finger domain, along with a region similar to the co-repressor protein N-Cor. MTA1 is normally expressed at low levels in various tissues and is more highly expressed in testis. Overexpression of MTA1 correlates with tumor invasion and metastasis in various carcinomas including colorectal, gastrointestinal and breast carcinomas. Elevation of MTA1 levels in these tumors appears to enhance the metastases to lymph nodes, increase mammary cell motility and potentiate growth, and therefore may be an indicator for assessing the potential malignancies of various tumors. A similar protein, MTA2, also designated MTA1-L1 (MTA1-like protein 1), shares more than 55 % sequence homology with MTA1 and is ubiquitously expressed.

---

Gene ID: 9219

Pathways: [Chromatin Binding](#)

## Application Details

---

Application Notes: WB 1:300-5000  
IHC-P 1:200-400  
IF(IHC-P) 1:50-200

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Concentration: 1 µg/µL

---

Buffer: 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

---

Preservative: ProClin

---

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

---

Storage: 4 °C, -20 °C

---

## Handling

---

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

---

Expiry Date: 12 months