antibodies -online.com





Datasheet for ABIN1387133

anti-BIN2 antibody



Overview

| Quantity: | 100 μL |
|--------------|--|
| Target: | BIN2 |
| Reactivity: | Human, Rat, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This BIN2 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human BIN2/BRAP1 |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Purified by Protein A. |

Target Details

| Target: | BIN2 |
|-------------------|--|
| Alternative Name: | BIN2 (BIN2 Products) |
| Background: | Synonyms: BIN2, BIN2_HUMAN, BRAP 1, Breast cancer associated protein BRAP1, Breast cancer-associated protein 1, Bridging integrator 2. |
| | Background: BAR proteins are characterized by a common N-terminal BAR (bin, amphiphysin |

and Rvs161/167) domain and are recognized as adaptor proteins that are involved in many cellular processes. BIN1 and BIN2 are BAR proteins that share 61 % sequence similarity. BIN1 (Bridging integrator 1) is a ubiquitously expressed regulatory protein for synaptic vesicle endocytosis. BIN1 also interacts with the transcription factors c-Myc and MyoD, potentially functioning as a tumor suppressor. BIN2, also known as Breast cancer-associated protein 1, is a 565 amino acid protein that interacts with BIN1. In contrast to BIN1, BIN2 lacks tumor suppressor features as well as a c-Myc interacting region. BIN2 shows preferred expression in tissues of hematopoietic origin, with high levels found in spleen, thymus, colon, placenta, lymphoid and granulocytic cells. There are two isoforms of BIN2 that are produced as a result of alternative splicing events.

Gene ID:

51411

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| Expiry Date: | 12 months |