# antibodies .- online.com





## Datasheet for ABIN1387151

### anti-ANKRD50 antibody

## Go to Product page

/ N	1	-	K	/1	01	A /
	1//	$\vdash$	1 \	/ I	e	<b>\/\/</b>
$\sim$	٧.	$\sim$	1 V		$\sim$	v v

Quantity:	100 μL	
Target:	ANKRD50	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ANKRD50 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ANKRD50
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

#### **Target Details**

Target:	ANKRD50
Alternative Name:	ANKRD50 (ANKRD50 Products)
Background:	Synonyms: ANKRD50, Ankyrin repeat domain 50, Ankyrin repeat domain-containing protein 50, ANR50_HUMAN, KIAA1223.
	Background: Ankyrins are membrane adaptor molecules that play important roles in coupling

integral membrane proteins to the spectrin-based cytoskeleton network. Mutations of ankyrin genes lead to severe genetic diseases, such as fatal cardiac arrhythmias and hereditary spherocytosis. ANKRD50 (ankyrin repeat domain 50) is a 1,429 amino acid phosphoprotein that contains nineteen ANK repeats. Conserved in chimpanzee, dog, cow, mouse, rat, chicken, zebrafish, fruit fly and mosquito, ANKRD50 is encoded by a gene that maps to human chromosome 4q28.1. Chromosome 4 represents approximately 6 % of the human genome and contains nearly 900 genes. Notably, the Huntingtin gene, which encodes an expanded glutamine tract in cases of Huntington's disease, is located on chromosome 4. FGFR-3 is also encoded by a gene that maps to human chromosome 4 and has been associated with thanatophoric dwarfism, achondroplasia, Muenke syndrome and bladder cancer. Chromosome 4 is also linked to Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.

Gene ID:

57182

#### **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IF(IHC-P) 1:50-200

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months