antibodies - online.com







anti-AFAP antibody (AA 286-325)



Image



Overview

Quantity:	100 μL
Target:	AFAP (AFAP1)
Binding Specificity:	AA 286-325
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AFAP antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human AFAP
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	AFAP (AFAP1)
Alternative Name:	AFAP (AFAP1 Products)
Background:	Synonyms: AFAP, AFAP110, AFAP-110, Actin filament-associated protein 1, 110 kDa actin

	filament-associated protein, AFAP1
	Background: Can cross-link actin filaments into both network and bundle structures (By
	similarity). May modulate changes in actin filament integrity and induce lamellipodia formation.
	May function as an adapter molecule that links other proteins, such as SRC and PKC to the
	actin cytoskeleton. Seems to play a role in the development and progression of prostate
	adenocarcinoma by regulating cell-matrix adhesions and migration in the cancer cells.
Gene ID:	60312

Application Details

UniProt:


Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

Restrictions: For Research Use only

Q8N556

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



Immunohistochemistry

Image 1. Formalin-fixed and paraffin embedded human lymphoma tissue labeled with Anti-AFAP/AFAP-110 Polyclonal Antibody, Unconjugated (ABIN1387171) at 1:200 followed by conjugation to the secondary antibody