# antibodies .- online.com







# anti-CCSAP antibody



( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	CCSAP
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCSAP antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human C1orf96
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

# **Target Details**

Target:	CCSAP	
Alternative Name:	C1orf96 (CCSAP Products)	
Background:	Synonyms: C1orf96, FLJ37296, FLJ41471, Uncharacterized protein C1orf96, CCSAP_HUMAN, Centriole, cilia and spindle-associated protein, CCSAP, CSAP, RP4-803J11.3.	
	Background: Chromosome 1 is the largest human chromosome spanning about 260 million	

base pairs and making up 8 % of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes lamin A. When defective, the LMNA gene product can build up in the nucleus and cause characteristic nuclear blebs. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma. The C1orf96 gene product has been provisionally designated C1orf96 pending further characterization.

Gene ID:

126731

## **Application Details**

Application Notes:

WB 1:300-5000

IHC-P 1:200-400

IF(IHC-P) 1:50-200

Restrictions:

For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months