# antibodies -online.com





## Datasheet for ABIN1387212

# anti-SH2D1A antibody



#### Overview

Quantity:	100 μL
Target:	SH2D1A
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SH2D1A antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human SH2D1A/SAP
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

### **Target Details**

SH2D1A
SH2D1A (SH2D1A Products)
Synonyms: SH2 domain protein 1A, DSHP, Duncan disease SH2 protein, SAP, Signaling lymphocyte activation molecule associated protein, SLAM associated protein, T cell signal transduction molecule SAP, SH21A_HUMAN.

Background: SH2D1A, also SH2 domain protein 1A, SAP and CD150/SLAM (signaling lymphocyte activation molecule)-associated protein, influences signaling pathways involving SLAM molecules at the interface between T and B cells. SH2D1A modulates SLAM by blocking the recruitment of tyrosine phosphatase SHP2 to the phosphorylated cytoplasmic domain of SLAM. SLAM activation mediates expansion of activated T cells during immune responses, induces production of interferon-?and changes the functional profile of subsets of T cells. SH2D1A is a hydrophilic, 128 amino acid protein that is 96 % homologous to the mouse protein in both SH2 and tail domains. SH2D1A is present in all major subsets of T cells, including CD4+, CD45RO+, CD45RA+ and CD8+, but not in B cells. SH2D1A can interact via an SH2 domain with a motif (TIYXXV) present in the cytoplasmic tail of cell-surface receptors SLAM (CD150), CD84, CD229 (LY9) and CD244 (2B4).

Gene ID:

4068

Pathways:

Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process

#### **Application Details**

Application Notes:

WB 1:300-5000

IHC-P 1:200-400

IF(IHC-P) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months