antibodies -online.com







anti-PCDH17 antibody (AA 31-130)



/ N	11/0	K /	\sim	A /
	ve	1 \/	ı⊢۱	Λ
\sim	' V C	1 V		v v

Quantity:	100 μL
Target:	PCDH17
Binding Specificity:	AA 31-130
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PCDH17 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PCDH17	
Isotype:	IgG	
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse	
Purification:	Purified by Protein A.	
Target Details		

Target:	PCDH17
Alternative Name:	PCDH17 (PCDH17 Products)

Target Details

Background:

Synonyms: PCDH68, C030033F14Rik, Gm78, PCD17_HUMAN, PCDH17, PCDH-17, PCDH 17, PCH68, Protocadherin 17, Protocadherin 68, Protocadherin-17, Protocadherin-68.

Background: Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin gene clusters, designated alpha, beta and gamma, all of which contain multiple tandemly arranged genes. PCDH17 is a 1,159 amino acid single-pass type I membrane protein that contains six cadherin domains. Expressed as multiple alternatively spliced isoforms, PCDH17 is thought to function as a calcium-dependent cell adhesion protein that may play a role in establishing cell-cell connections within brain tissue. The gene encoding PCDH17 maps to human chromosome 13, which houses over 400 genes, such as BRCA2 and RB1, and comprises nearly 4 % of the human genome.

Application Details

Application Notes:	ELISA 1:500-1000	
	IHC-P 1:200-400	

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

ICC 1:100-500

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months