antibodies -online.com





Datasheet for ABIN1387288

anti-KCNAB1 antibody (AA 131-230)



Overview

Quantity:	100 μL
Target:	KCNAB1
Binding Specificity:	AA 131-230
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KCNAB1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human KCNAB1/Kv beta 1
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Pig,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target: KCNAB1

Target Details

Alternative Name:	KCNAB1 (KCNAB1 Products)	
Background:	Synonyms: hKvb3; hKvBeta3; K+ channel subunit beta-1; KCAB1_HUMAN; KCNA1B; KCNAB1;	
	KV-BETA-1; Kvb1.3; Voltage-gated potassium channel beta-1 subunit; Voltage-gated potassium	
	channel subunit beta-1.	
	Background: Voltage-gated K+ channels in the plasma membrane control the repolarization an	
	the frequency of action potentials in neurons, muscles and other excitable cells. The KV gene	
	family encodes more than 30 proteins that comprise the subunits of the K+ channels, and they	
	vary in their gating and permeation properties, subcellular distribution and expression patterns.	
	Functional KV channels assemble as tetramers consisting of pore-forming å subunits (KV),	
	which include the KV1, KV2, KV3 and KV4 proteins, and accessory or KV-subunits that modify	
	the gating properties of the coexpressed KV subunits. KV, also known as KCNAB1 (potassium	
	voltage-gated channel, shaker-related subfamily, beta member 1), is a 419 amino acid	
	accessory K+ channel protein that exists as three alternatively spliced isoforms and regulates	
	the activity of the pore-forming å subunit. It is expressed in brain, with highest levels detected in	
	caudate nucleus, hippocampus and thalamus.	
Gene ID:	7881	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months