# antibodies - online.com





Datasheet for ABIN1387302

## anti-LGI3 antibody (AA 101-220)



#### Overview

Quantity:	100 μL
Target:	LGI3
Binding Specificity:	AA 101-220
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LGI3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human LGI3
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Sheep,Pig,Horse
Purification:	Purified by Protein A.

#### **Target Details**

et:
-----

## Target Details

Precaution of Use:

round: S	Synonyms: Leucine rich repeat LGI family member 3; Leucine-rich glioma inactivated protein 3;
L	Leucine-rich glioma-inactivated protein 3; Leucine-rich repeat LGI family member 3; LGI1 like
р	protein 4; LGI1-like protein 4; LGI3; LGI3_HUMAN; LGIL4.
В	Background: The leucine-rich (LRR) repeat is a 20-30 amino acid motif that forms a
h	nydrophobic å/ horseshoe fold, allowing it to accommodate several leucine residues within a
ti	ightly packed core. All LRR repeats contain a variable segment and a highly conserved
S	segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. Leucine-rich
g	glioma-inactivated protein 3 (LGI3), also known as LGI1-like protein 4 (LGIL4) or leucine-rich
re	repeat LGI family member 3, is a 548 amino acid secretory protein. LGI3 contains five LRR
re	repeats and seven EAR repeats. Widely expressed, with highest levels in brain and lung, LGI3
h	has been shown to colocalize with endocytosis-associated proteins, lipid raft markers, and
S	Syntaxin. The gene encoding LGI3 maps to chromosome 8p21.3.
cation Details	
ation Notes: V	WB 1:300-5000
E	ELISA 1:500-1000
II	HC-P 1:200-400
II	HC-F 1:100-500
II	F(IHC-P) 1:50-200
11	F(IHC-F) 1:50-200
11	F(ICC) 1:50-200
I	CC 1:100-500
ctions: F	For Research Use only
lling	
ıt: L	_iquid
ntration: 1	1 μg/μL
: 0	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
vative: P	ProClin
etions: F  Hing  It: L  Intration: 1	F(ICC) 1:50-200 CC 1:100-500 For Research Use only Liquid 1 µg/µL 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

handled by trained staff only.

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

### Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months