antibodies -online.com







anti-MSH3 antibody (AA 651-750)



Image



Overview

Quantity:	100 μL
Target:	MSH3
Binding Specificity:	AA 651-750
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MSH3 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Target:

Alternative Name:

1 Toddet Details	
Immunogen:	KLH conjugated synthetic peptide derived from human MSH3
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Rabbit
Purification:	Purified by Protein A.
Target Details	

MSH3

MSH3 (MSH3 Products)

Target Details

Storage Comment:


rarget Details	
Background:	Synonyms: Divergent upstream protein, DNA mismatch repair protein, DNA mismatch repair
	protein Msh 3, DNA mismatch repair protein Msh3, DUC 1, DUC1, DUG, DUP, hMSH3, Mismatch
	repair protein 1, MRP 1, MRP1, MSH 3, MSH-3, MSH3_HUMAN, MutS homolog 3.
	Background: Component of the post-replicative DNA mismatch repair system (MMR).
	Heterodimerizes with MSH2 to form MutS beta which binds to DNA mismatches thereby
	initiating DNA repair. When bound, the MutS beta heterodimer bends the DNA helix and shields
	approximately 20 base pairs. MutS beta recognizes large insertion-deletion loops (IDL) up to 13
	nucleotides long. After mismatch binding, forms a ternary complex with the MutL alpha
	heterodimer, which is thought to be responsible for directing the downstream MMR events,
	including strand discrimination, excision, and resynthesis.
Gene ID:	4437
Pathways:	DNA Damage Repair, Production of Molecular Mediator of Immune Response
Application Details	
Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date:

12 months

Images



ELISA

lmage 1. Antigen: 2 μ g/100 μ L Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000; Secondary: HRP conjugated Rabbit Anti-Goat IgG at 1: 5000; TMB staining Read the data in Microplate Reader by 450nm.