antibodies -online.com





anti-FOXG1 antibody (AA 201-300)



Overview

Quantity:	100 μL
Target:	FOXG1
Binding Specificity:	AA 201-300
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FOXG1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FOXG1
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	FOXG1
Alternative Name:	FOXG1/BF-1 (FOXG1 Products)
Background:	Synonyms: BF 1, BF 2, BF-1, BF-2, BF1, BF2 , Brain factor 1, Brain factor 2, FHKL, FKH2, FKHL1,

FKHL2, FKHL3, FKHL4, Forkhead box G1A, Forkhead box G1B, Forkhead box protein G1, Forkhead box protein G1A, Forkhead box protein G1B, Forkhead box protein G1C, Forkhead drosophila homolog like 2, Forkhead like 1, Forkhead like 2, FOXG1_HUMAN.

Background: The winged-helix transcriptional repressor (WH) BF-1 gene encodes brain factor 1 (BF-1), also known as foxg1, and is essential for the proliferation of progenitor cells in the cerebral cortex and influences regional patterning in the mammalian telencephalon (14). WH proteins are a family of putative transcriptional regulators with diverse roles in development, and are characterized by a highly conserved DNA binding structure, the WH domain (1,5,6). BF-1 plays a critical role in the development of the cerebral hemispheres of the brain and targeted disruption of the gene leads to severe defects in the development of telencephalic structures, such as the cerebral cortex and basal ganglia (1). The loss of BF-1 results in an accelerated rate of neuronal differentiation and the shortening of the neurogenetic period in the embryonic cerebral cortex (1,7). BF-1 is expressed by E8.5 in telencephalic progenitors (1). It may also regulate the response of cerebral cortical progenitors to environmental cues (1).

Application Details

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months