# antibodies -online.com







# anti-Advillin antibody (AA 151-250)



#### Overview

Quantity:	100 μL
Target:	Advillin (AVIL)
Binding Specificity:	AA 151-250
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Advillin antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Advillin
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

#### **Target Details**

Target: Advillin (AVIL)

## Target Details

Alternative Name:	Advillia (AVIII Droducto)
Alternative Name:	Advillin (AVIL Products)
Background:	Synonyms: Actin binding protein DOC 6, Actin binding protein DOC6, Advil, Advillin, AVIL,
	AVIL_HUMAN, DOC 6, DOC6, p92.
	Background: Advillin is an 819 amino acid protein that localizes to both the cytoplasm and the
	cytoskeleton and contains one HP domain and six gelsolin-like repeats. Expressed at high level
	in colon and small intestine and at lower levels in uterus, thymus, testis and prostate, advillin
	functions as a calcium-regulated Actin-binding protein that may be involved in the development
	of neuronal cells, specifically those that form ganglia. The gene encoding advillin maps to
	human chromosome 12, which encodes over 1,100 genes and comprises approximately $4.5\%$
	of the human genome. Chromosome 12 is associated with a variety of diseases and afflictions
	including hypochondrogenesis, achondrogenesis, Kniest dysplasia, Noonan syndrome and
	Trisomy 12p, which causes facial developmental defects and seizure disorders.
Gene ID:	3316
Pathways:	Regulation of Actin Filament Polymerization
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

## Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months