

Datasheet for ABIN1387572  
**anti-H13 antibody (AA 251-350)**[Go to Product page](#)

## 2 Images

## Overview

|                      |   |
|----------------------|---|
| Quantity:            | 100 µL  |
| Target:              | H13   |
| Binding Specificity: | AA 251-350  |
| Reactivity:          | Human   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This H13 antibody is un-conjugated  |
| Application:         | Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)),<br>Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-<br>embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)),<br>Immunocytochemistry (ICC) |

## Product Details

|                       |  |
|-----------------------|--|
| Immunogen:            | KLH conjugated synthetic peptide derived from human Signal Peptide Peptidase |
| Isotype:              | IgG  |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit                             |
| Purification:         | Purified by Protein A.   |

## Target Details

|                   |   |
|-------------------|---|
| Target:           | H13   |
| Alternative Name: | Spp/Signal Peptide Peptidase ( <a href="#">H13 Products</a> ) |

## Target Details

|             |  |
|-------------|--|
| Background: | <p>Synonyms: Histocompatibility minor 13, HM 13, HM13, HM13_HUMAN, IMP 1, IMP-1, IMP1, IMPAS, IMPAS-1, Intramembrane Protease 1, Intramembrane protease, Minor Histocompatibility 13, Minor histocompatibility antigen 13, Minor histocompatibility antigen H13, MSTP086, Presenilin like protein 3, Presenilin-like protein 3, PSEN1 3, PSEN13, PSL 3, PSL3, Signal peptide peptidase, Signal peptide peptidase beta, SPP, dJ324017.1, H13, hIMP1, hIMP1 protein.</p> <p>Background: The endoplasmic reticulum exerts a quality control over newly synthesized proteins and a variety of components have been implicated in the specific recognition of aberrant or misfolded polypeptides. Signal peptide peptidase (SPP) catalyzes intramembrane proteolysis of some signal peptides after they have been cleaved from a preprotein, resulting in the release of the fragment from the ER membrane into the cytoplasm. SPP is required to generate lymphocyte cell surface (HLA-E) epitopes derived from MHC class I signal peptides, and may play a role in graft rejection. It also may be necessary for the removal of the signal peptide that remains attached to the hepatitis C virus core protein after the initial proteolytic processing of the polyprotein.</p> |
|-------------|--|

|          |     |
|----------|-----|
| Gene ID: | 207 |
|----------|-----|

## Application Details

|                    |  |
|--------------------|--|
| Application Notes: | WB 1:300-5000<br>ELISA 1:500-1000<br>IHC-P 1:200-400<br>IHC-F 1:100-500<br>IF(IHC-P) 1:50-200<br>IF(IHC-F) 1:50-200<br>IF(ICC) 1:50-200<br>ICC 1:100-500 |
|--------------------|--|

|               |                       |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|


## Handling

|                |   |
|----------------|---|
| Format:        | Liquid  |
| Concentration: | 1 µg/µL   |
| Buffer:        | 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. |
| Preservative:  | ProClin   |

Handling


|                    |  |
|--------------------|--|
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | 4 °C,-20 °C  |
| Storage Comment:   | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.                                    |
| Expiry Date:       | 12 months  |

Images



**ELISA**

**Image 1.** Antigen: 2 µg/100 µL Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000; Secondary: HRP conjugated Rabbit Anti-Goat IgG at 1: 5000; TMB staining Read the data in Microplate Reader by 450nm.



**ELISA**

**Image 2.** Antigen: 0.2ug/100ul, Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000, Secondary: HRP conjugated Goat-Anti-Rabbit IgG at 1: 5000, TMB staining, Read the data in MicroplateReader by 450nm