# antibodies -online.com





anti-RND2 antibody (AA 1-80)



#### Overview

Quantity:	100 μL
Target:	RND2
Binding Specificity:	AA 1-80
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RND2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human RND2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

### **Target Details**

|--|--|

## Target Details

Alternative Name:	RND2/Rho7 (RND2 Products)
Background:	Synonyms: ARHN, Rho 7, Rho family GTPase 2, Rho N, Rho related GTP-binding protein RhoN,
	Rho-related GTP-binding protein Rho7, Rho-related GTP-binding protein RhoN, Rho7, RhoN, Rno
	2, Rnd2, RND2_HUMAN.
	Background: The Ras p21 family of guanine nucleotide proteins has been widely studied in view
	of its apparent role in signal transduction pathways and high frequency of mutations in human
	malignancies. It is now clear, however, that the Ras proteins (H-, K- and N-Ras p21) are
	members of a much larger superfamily of related proteins. Six members of this family, Rap 1A,
	Rap 1B, Rap 2, R-Ras, Ral A and Ral B, exhibit approximately 50 % amino acid homology to Ras.
	The six mammalian Rho proteins (Rho A, B, C, G, 7 and 8) are approximately 30 % homologous
	to Ras and are expressed in a wide range of cell types. Both Ras p21 and Rho p21, as well as
	other members of the Ras superfamily, contain a carboxy-terminal CAAX sequence (C, cysteine
	A, aliphatic amino acid, X, any amino acid) which in the case of Ras has been shown to be
	essential for correct localization and function.
Pathways:	Regulation of Cell Size
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

## Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months