

[Go to Product page](#)

Datasheet for ABIN1387670 **anti-KCTD12 antibody (AA 251-325)**

Overview

Quantity:	100 µL
Target:	KCTD12
Binding Specificity:	AA 251-325
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KCTD12 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human KCTD12
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	KCTD12
Alternative Name:	KCTD12 (KCTD12 Products)

Target Details

Background:	<p>Synonyms: C13orf2, BTB/POZ domain-containing protein KCTD12, C13orf2, KCD12_HUMAN, KCTD12, KIAA1778, PFET1, Pfetin, potassium channel tetramerisation domain containing 12 USA, Predominantly fetal expressed T1 domain.</p> <p>Background: The BTB (Broad-Complex, Tramtrack and Bric a brac) domain, also known as the POZ (POxvirus and Zinc finger) domain, is an N-terminal homodimerization domain that contains multiple copies of kelch repeats and/or C2H2-type zinc fingers. Proteins that contain BTB domains are thought to be involved in transcriptional regulation via control of chromatin structure and function. KCTD12 (potassium channel tetramerisation domain containing 12), also known as PFET1 or PFETIN, is a 325 amino acid protein that is expressed in fetal organs, with highest levels in the cochlea and brain and extremely low levels in adult organs, such as brain and lung. KCTD12 is considered a prognostic biomarker of gastrointestinal stromal tumors.</p>
-------------	---

Pathways:	Synaptic Membrane , Regulation of G-Protein Coupled Receptor Protein Signaling
-----------	--

Application Details

Application Notes:	ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C, -20 °C

Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months