Datasheet for ABIN1387743

# anti-C22orf28 antibody

Go to Product page

## Overview

| | |
|---|---|
| Quantity: | 100 µL |
| Target: | C22orf28 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This C22orf28 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

## Product Details

| | |
|---|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human HSPC117/C22orf28 |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Purified by Protein A. |

## Target Details

| | |
|---|---|
| Target: | C22orf28 |
| Alternative Name: | C22orf28 (C22orf28 Products) |
| Background: | Synonyms: FAAP, Ankyrin repeat domain 54, C22orf28, Chromosome 22 open reading frame 28, DJ149A16.6, HSPC117, Hypothetical protein LOC51493, Novel protein HSPC117, RP1-149A16.6, RTCB_HUMAN, tRNA-splicing ligase RtcB homolog. |

Copyright antibodies-online. All rights reserved.

## Target Details

| | |
|---|---|
| | Background: HSPC117, also known as C22orf28, is a 505 amino acid protein that is encoded by a gene which maps to human chromosome 22. A highly homologous protein identified in rodents, FAAP (focal adhesion associated protein), encoded by murine D10Wsu52e gene, has been suggested to play a role in regulating cell adhesion dynamics. Chromosome 22 houses over 500 genes and is the second smallest human chromosome. Mutations in several of the genes that map to chromosome 22 are involved in the development of Phelan-McDermid syndrome, Neurofibromatosis type 2, autism and schizophrenia. Additionally, translocations between chromosomes 9 and 22 may lead to the formation of the Philadelphia Chromosome and the subsequent production of the novel fusion protein BCR-Abl, a potent cell proliferation activator found in several types of leukemias. |
| Gene ID: | 51493 |

## Application Details

| | |
|---|---|
| Application Notes: | WB 1:300-5000<br>IHC-P 1:200-400<br>IF(IHC-P) 1:50-200 |
| Restrictions: | For Research Use only |

## Handling

| | |
|---|---|
| Format: | Liquid |
| Concentration: | 1 µg/µL |
| Buffer: | 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| Expiry Date: | 12 months |

 Copyright antibodies-online. All rights reserved.