antibodies -online.com





anti-Semaphorin 6C antibody (AA 256-305)



Go to Product page

\sim			
	N/F	ا/\r14	$\triangle W$

Quantity:	100 μL
Target:	Semaphorin 6C (SEMA6C)
Binding Specificity:	AA 256-305
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Semaphorin 6C antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SEMA6C
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Semaphorin 6C (SEMA6C)
Alternative Name:	SEMA6C (SEMA6C Products)

Target Details

Background:

Synonyms: KIAA1869, m Sema Y, m Sema Y2, sema domain, transmembrane domain TM, and cytoplasmic domain, semaphorin 6C, Sema Y antibody Semaphorin 6C [Precursor], Semaphorin Y, SEMAY, SEM6C_HUMAN.

Background: Semaphorins are a family of cell surface and secreted proteins involved in neural development that are conserved from insects to humans. Members of this family are approximately 750 amino acids in length (including signal sequences) and are defined by a conserved extracellular semaphorin? domain of approximately 500 amino acids containing 14-16 cysteines, blocks of conserved sequences and no obvious repeats. The transmembrane semaphorins are characterized by an additional 80 amino acid transmembrane domain and an 80-110 amino acid cytoplasmic domain. SEMA6C, also known as SEMA Y, is a transmembrane protein expressed in fetal brain and adult skeletal muscle. Three isoforms of this semaphorin exist due to alternative splicing: SEMA6C 1, SEMA6C 2 and SEMA6C 3. The extracellular domain of SEMA6C induces growth cone collapse of dorsal root ganglion and plays a role in generation or stability of entorhino-hippocampal synapses.

Application Details

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months