# antibodies .- online.com





Datasheet for ABIN1387766

## anti-SOX6 antibody (AA 1-100)



#### Overview

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | SOX6  |
| Binding Specificity: | AA 1-100  |
| Reactivity:          | Human   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This SOX6 antibody is un-conjugated   |
| Application:         | ELISA, Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human SOX6 |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Human  |
| Predicted Reactivity: | Mouse,Rat,Dog,Cow,Sheep,Pig,Rabbit                       |
| Purification:         | Purified by Protein A.                                   |

#### **Target Details**

Target: SOX6

### **Target Details**

| Alternative Name:   | SOX6 (SOX6 Products)   |  |
|---------------------|--|--|
| Background:         | Synonyms: HSSOX 6, HSSOX6, SOX-6 SOX 6, SOX6_HUMAN.  |  |
|                     | Background: Sox genes comprise a family of genes that are related to the mammalian sex                             |  |
|                     | determining gene SRY. These genes similarly contain sequences that encode for the HMG-box                          |  |
|                     | domain, which is responsible for the sequence-specific DNA-binding activity. Sox genes encode                      |  |
|                     | putative transcriptional regulators implicated in the decision of cell fates during development                    |  |
|                     | and the control of diverse developmental processes. The highly complex group of Sox genes                          |  |
|                     | cluster at least 40 different loci that rapidly diverged in various animal lineages. At present, 30                |  |
|                     | Sox genes have been identified. Members of this family have been shown to be conserved                             |  |
|                     | during evolution and to play key roles during animal development. Some are involved in human                       |  |
|                     | diseases, including sex reversal.  |  |
| Application Details |  |  |
| Application Notes:  | ELISA 1:500-1000   |  |
|                     | FCM 1:20-100   |  |
|                     | IHC-P 1:200-400  |  |
|                     | IHC-F 1:100-500  |  |
|                     | IF(IHC-P) 1:50-200   |  |
|                     | IF(IHC-F) 1:50-200   |  |
|                     | IF(ICC) 1:50-200   |  |
|                     | ICC 1:100-500  |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Concentration:      | 1 μg/μL  |  |
| Buffer:             | 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.  |  |
| Preservative:       | ProClin  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |  |
| Storage:            | 4 °C,-20 °C  |  |
| Storage Comment:    | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.                                    |  |
|                     |  |  |

|          |       | 1.   |   |
|----------|-------|------|---|
| $\vdash$ | land  | lın  |   |
|          | ıaııu | 1111 | U |
|          |       |      |   |

Expiry Date:

12 months