antibodies -online.com





anti-RAPGEF3 antibody (AA 301-400)

2 Images



Overview

Quantity:	100 μL
Target:	RAPGEF3
Binding Specificity:	AA 301-400
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RAPGEF3 antibody is un-conjugated
Application:	ELISA, Immunocytochemistry (ICC), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Epac1
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Rabbit
Purification:	Purified by Protein A.

Target Details

|--|


Target Details

Alternative Name:	Epac1 (RAPGEF3 Products)
Background:	Synonyms: bcm910, CAMP GEFI, cAMP regulated guanine nucleotide exchange factor I,
	CAMPGEFI, CGEF 1, CGEF1, EPA1, Epac 1, EPAC, EPAC1, Exchange factor directly activated by
	cAMP 1, Exchange protein directly activated by cAMP 1, MGC21410, RAP guanine nucleotide
	exchange factor, Rap guanine nucleotide exchange factor GEF 3, RAP guanine nucleotide
	exchange factor 3, Rap1 guanine nucleotide exchange factor directly activated by cAMP,
	RAPGEF3.
	Background: The activation of RaP1 by cAMP is independent of PKA and is mediated by
	recently discovered family of guanine nucleotide exchange factors (GEFs) called cAMP-GEFs of
	Epacs. The Epac signaling therefore represents a novel mechanism for cAMP signaling with in
	the cAMP cascade. There are 2 members of the Epac family, Epac1 and Epac 2. Both proteins
	are multidomain proteins containing an autoinhibitory cAMP-binding domain that inhibits the
	catalytic region and a DEP domain (dishevelled, Egl-10 and pleckstrin homology domain)
	targeting the membrane anchors. EPAC2 has an additional cAMP-binding site in its N-terminus
	that binds cAMP with low affinity. EPAC1 mRNA is broadly expressed, with particularly high
	levels occurring in the thyroid, ovary, kidney and certain brain regions, whereas expression of
	EPAC2 mRNA appears to be restricted to the brain and adrenal glands. Epac 1 and Epac 2 also
	interact with light chain 2 (LC2) or MAP1A that serves as a scaffolding structure to stabilize the
	signal transduction complex. The Epac 1-selective were generated against unique antigenic
	sequences form near N-terminus and between RasGEFN and Ras GEF domains. The to Epac
	1are affinity purified over immobilized antigen based chromatography.
Gene ID:	1019
Pathways:	cAMP Metabolic Process
Application Details	
Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only

Handling


Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Images



ELISA

Image 1. Antigen: 0.2 μ g/100 μ L Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000; Secondary: HRP conjugated Goat-Anti-Rabbit IgG at 1: 5000; TMB staining; Read the data in MicroplateReader by 450



ELISA

Image 2. Antigen: 0.2ug/100ul, Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000, Secondary: HRP conjugated Goat-Anti-Rabbit IgG at 1: 5000, TMB staining, Read the data in MicroplateReader by 450nm