antibodies -online.com







anti-Netrin 4 antibody (AA 520-628) (Biotin)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	Netrin 4 (NTN4)	
Binding Specificity:	AA 520-628	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Netrin 4 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Netrin 4	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	Netrin 4 (NTN4)	
Alternative Name:	Netrin 4 (NTN4 Products)	
Background:	kground: Synonyms: Beta netrin, Beta-netrin, Hepar derived netrin like protein, Hepar-derived netrin-li	

protein, NET4_HUMAN, Netrin-4, Netrin-4, NTN 4, NTN4.

Background: Netrin proteins are a family of laminin-related secreted proteins that provide guidance signals for axonal growth and cell migration during development. Netrin signaling is dependent on the concentration of calcium outside the cell and the level of PKA activity. In axonal cells, a reduction in PKA activity converts the responsiveness of the axons to the netrin proteins, as the cells are repelled, rather than attracted, by the netrin gradient. Netrin-4 is related to the Laminin chains, and is therefore also designated -netrin. It is present in the basement membranes of the vasculature, lateral olfactory tract, kidney and ovary. In humans, the gene encoding for the netrin-4 protein is localized to chromosome 12q22-q23. High levels of netrin-4 mRNA have also been detected in many cells and tissues, including cerebral cortex, hippocampus, amygdaloid nuclei and Purkinje cells. Netrin-4 is important in neural, kidney and vascular development.

Gene ID:

59277

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months