antibodies -online.com





anti-ACADL antibody (AA 31-130) (Biotin)



Go to Product page

()	11/0	K\ /	iew
	\cup	ועוי	$I \cap VV$

Quantity:	100 μL
Target:	ACADL
Binding Specificity:	AA 31-130
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ACADL antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ACADL
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat
Purification:	Purified by Protein A.

Target Details

Target:	ACADL
Alternative Name:	ACADL (ACADL Products)
Background:	Synonyms: mitochondrial, ACAD4, ACADL, ACADL_HUMAN, Acyl Coenzyme A dehydrogenase

long chain, FLJ94052, LCAD, Long chain acyl CoA dehydrogenase, Long-chain specic acyl-CoA dehydrogenase.

Background: The protein encoded by this gene belongs to the acyl-CoA dehydrogenase family, which is a family of mitochondrial flavoenzymes involved in fatty acid and branched chain amino-acid metabolism. This protein is one of the four enzymes that catalyze the initial step of mitochondrial beta-oxidation of straight-chain fatty acid. Defects in this gene are the cause of long-chain acyl-CoA dehydrogenase (LCAD) deficiency, leading to nonketotic hypoglycemia. [provided by RefSeq].

Pathways:

Monocarboxylic Acid Catabolic Process

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400 IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Storage:

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage Comment: Store at -20°C for 12 months.

-20 °C

Expiry Date: 12 months