antibodies - online.com





anti-CRIPT antibody (AA 21-80) (Biotin)



	_				
Go t	\cap \vdash	rod	uct	nac	пe

_							
0	V	е	r١	/1	е	V	1

Quantity:	100 μL	
Target:	CRIPT	
Binding Specificity:	AA 21-80	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CRIPT antibody is conjugated to Biotin	
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CRIPT	
Isotype:	IgG	
Predicted Reactivity: Human, Mouse, Rat, Cow, Pig, Chicken		
Purification:	Purified by Protein A.	

Target Details

Target:	CRIPT	
Alternative Name:	CRIPT (CRIPT Products)	
Background: Synonyms: CRIPT, CRIPT_HUMAN, Cysteine rich interactor of PDZ3, Cysteine		

protein, Cysteine-rich interactor of PDZ three, Cysteine-rich interactor of PDZ3, Cysteine-rich PDZ-binding protein, HSPC139, HSPC139 protein, Hypothetical protein CRIPT, Postsynaptic protein CRIPT.

Background: The PSD-95/SAP 90 family of proteins, which are known to bind to and cluster various membrane proteins, are involved in the organization of synaptic structure. These proteins are physically and functionally linked to cytoskeletal and/or signaling proteins. CRIPT (for cysteine-rich interactor of PDZ three), a novel postsynaptic protein, binds specifically to the PDZ3 domain of PSD-95/SAP 90. CRIPT induces the recruitment of PSD-95/SAP 90 to microtubules, and it has been shown to bind directly to microtubules, indicating that it may be responsible for cytoskeletal anchoring of PSD-95/SAP 90. CRIPT is widely expressed outside of the brain and is highly conserved from animals to plants suggesting a wider role in regulating cytoskeleton-membrane associations.

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months