antibodies -online.com





anti-MAP1B antibody (AA 451-550) (Alexa Fluor 350)



| () | 1 / | \sim | KI / | 110 | Νę |
|-----|-----|----------|------|-----|-------|
| | 1// | \vdash | I \/ | 1 ← | ٠// ٢ |
| | | | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | MAP1B | |
| Binding Specificity: | AA 451-550 | |
| Reactivity: | Human, Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This MAP1B antibody is conjugated to Alexa Fluor 350 | |
| Application: | Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human MAP1B | |
|-----------------------|---|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse, Rat | |
| Predicted Reactivity: | Dog,Cow,Sheep,Pig,Horse,Rabbit | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | MAP1B |
|-------------------|------------------------|
| Alternative Name: | MAP1B (MAP1B Products) |

Target Details

| _ | | | |
|---------------------|---|--|--|
| Background: | Synonyms: MAP5, FUTSCH, PPP1R102, Microtubule-associated protein 1B, MAP-1B, MAP1B | | |
| | Background: Facilitates tyrosination of alpha-tubulin in neuronal microtubules (By similarity). | | |
| | Phosphorylated MAP1B may play a role in the cytoskeletal changes that accompany neurite | | |
| | extension. Possibly MAP1B binds to at least two tubulin subunits in the polymer, and this | | |
| | bridging of subunits might be involved in nucleating microtubule polymerization and in | | |
| | stabilizing microtubules. Acts as a positive cofactor in DAPK1-mediated autophagic vesicle | | |
| | formation and membrane blebbing. | | |
| Gene ID: | 4131 | | |
| UniProt: | P46821 | | |
| Pathways: | Microtubule Dynamics, Regulation of Cell Size | | |
| Application Details | | | |
| Application Notes: | FCM 1:20-100 | | |
| | IF(IHC-P) 1:50-200 | | |
| | IF(IHC-F) 1:50-200 | | |
| | IF(ICC) 1:50-200 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 1 μg/μL | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | | |
| | 50 % Glycerol. | | |
| Preservative: | ProClin | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | | |
| | handled by trained staff only. | | |
| Storage: | -20 °C | | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | | |
| Expiry Date: | 12 months | | |
| | | | |