

[Go to Product page](#)

Datasheet for ABIN1388849

anti-Midline 2 antibody (AA 251-350) (Alexa Fluor 555)

Overview

Quantity:	100 µL
Target:	Midline 2 (MID2)
Binding Specificity:	AA 251-350
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Midline 2 antibody is conjugated to Alexa Fluor 555
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RNF60/Midline-2
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Sheep,Pig,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	Midline 2 (MID2)
Alternative Name:	RNF60/Midline-2 (MID2 Products)

Target Details

Background:	<p>Synonyms: FXY2, MID2, Midin-2, Midline 2, Midline defect 2, Midline-2, Probable E3 ubiquitin-protein ligase MID2, RING finger protein 60, Ring finger protein protein 60, RNF60, TRIM1, TRIM1_HUMAN, Tripartite mot protein 1, Tripartite mot-containing protein 1, Tripartite mot-containing protein 1, Midline defect 2, FLJ37715, FLJ41813.</p> <p>Background: Midline-2 is a 715 amino acid protein encoded by the human gene MID2. Midline-2 belongs to the TRIM/RBCC family and contains two B box-type zinc fingers, one B30.2/SPRY domain, one COS domain, one fibronectin type-III domain and one RING-type zinc finger.</p> <p>Midline-2 is a cytoplasmic protein found as a homodimer or heterodimer with Midline-1. It also interacts with IGBP1 (Lymphocyte signaling protein A4). Dimerization is mediated by the tripartite motif (RBCC, RING- and B box-type zinc fingers and coiled coil domains) and microtubule association is dependent on the C-terminal B30.2 domain. Midline-2 is expressed at low levels in fetal kidney and lung, and in adult prostate, ovary and small intestine.</p>
-------------	---

Gene ID:	26040
----------	-------

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months