



[Go to Product page](#)

Datasheet for ABIN1388912  
**anti-STXBP6 antibody (AA 1-100) (FITC)**

### Overview

|                      |  |
|----------------------|--|
| Quantity:            | 100 µL   |
| Target:              | STXBP6   |
| Binding Specificity: | AA 1-100   |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This STXBP6 antibody is conjugated to FITC   |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

### Product Details

|                       |  |
|-----------------------|--|
| Immunogen:            | KLH conjugated synthetic peptide derived from human Amisyn |
| Isotype:              | IgG  |
| Predicted Reactivity: | Human, Mouse, Rat, Cow, Sheep, Chicken, Rabbit             |
| Purification:         | Purified by Protein A.                                     |

### Target Details

|                   |   |
|-------------------|---|
| Target:           | STXBP6  |
| Alternative Name: | Amisyn ( <a href="#">STXBP6 Products</a> )  |
| Background:       | Synonyms: Amisyn, HSPC156, STXB6_HUMAN, STXBP6, Syntaxin binding protein 6, Syntaxin- |

## Target Details

---

binding protein 6.

Background: Amisyn is a mostly cytosolic protein related to Tomosyn which plays an important role in SNARE complex assembly. Amisyn contains a v-SNARE coiled coil homology domain that binds to Syntaxin 1A and weakly to Syntaxin 4. Three isoforms exist for Amisyn. Isoform 1 is the full length protein, isoform 2 has a different amino acid sequence between residues 204-210 and isoform 3 is missing amino acids 1-102 and contains a different sequence for amino acids 103-150. Amisyn lacks a transmembrane domain and therefore is unable to assemble into a functional, membrane-anchored SNARE complex. This suggests that Amisyn may instead be acting to maintain SNARE conformation and facilitate the binding of VAMP-2. Amisyn can inhibit exocytosis independent of Syntaxin binding.

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months