



[Go to Product page](#)

Datasheet for ABIN1388917 **anti-Bestrophin antibody (Biotin)**

Overview

Quantity:	100 µL
Target:	Bestrophin (LOC100368263)
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Bestrophin antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Bestrophin
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Bestrophin (LOC100368263)
Alternative Name:	Bestrophin (LOC100368263 Products)
Background:	Synonyms: BEST 1, BEST1, BEST-1, BEST, Best macular dystrophy, BEST1, BEST1_HUMAN,

Target Details

Bestrophin 1, Bestrophin-1, Bestrophin1, BMD, mBest1, TU15B, Vitellorm macular dystrophy 2, Vitellorm macular dystrophy, Vitellorm macular dystrophy protein 2, VMD 2, VMD2.

Background: Best vitelliform macular dystrophy, known as Best disease, is an early-onset autosomal dominant condition in which accumulation of lipofuscin-like material within and beneath the RPE leads to progressive loss of central vision. Best disease is frequently a reflection of mutations in the Bestrophin gene, which encodes a protein containing four putative transmembrane domains and localizes to the basolateral plasma membrane of RPE cells. Human Bestrophin forms oligomeric chloride channels that are sensitive to intracellular calcium. Missense mutations at the Bestrophin locus reduces or abolishes Bestrophin protein mediated membrane current. Bestrophin 2, Bestrophin 3, and Bestrophin 4 are transmembrane proteins that contain a high percentage of aromatic residues, a conserved RFP (Arg-Phe-Pro) motif and they function as anion channels.

Gene ID: 5068

Application Details

Application Notes: WB 1:100-1000
IHC-P 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months