# antibodies -online.com







# anti-HAPLN2 antibody (AA 21-120) (Biotin)



| ( ) | 11/0   | K\ /   | iew                 | 1 |
|-----|--------|--------|---------------------|---|
|     | $\cup$ | 'I V/I | $\square \vee \vee$ | ı |
|     |        |        |                     |   |

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | HAPLN2  |
| Binding Specificity: | AA 21-120   |
| Reactivity:          | Human   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This HAPLN2 antibody is conjugated to Biotin  |
| Application:         | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

## **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human HAPLN2 |
|-----------------------|--|
| Isotype:              | IgG  |
| Predicted Reactivity: | Human,Mouse,Rat,Cow,Sheep,Pig,Rabbit                       |
| Purification:         | Purified by Protein A.                                     |

# **Target Details**

| Target:           | HAPLN2  |
|-------------------|---|
| Alternative Name: | HAPLN2 (HAPLN2 Products)  |
| Background:       | Synonyms: Brain link protein 1, Bral1, Hapln2, HPLN2_HUMAN, Hyaluronan and proteoglycan |

link protein 2.

Background: HAPLN2 is a 340 amino acid protein encoded by the human gene HAPLN2. HAPLN2 belongs to the HAPLN family and contains one immunoglobulin (Ig)-like, V-type domain and two link domains. HAPLN2 mediates a firm binding of versican V2 to hyaluronic acid. HAPLN2 is believed to play a pivotal role in the formation of the hyaluronan-associated matrix in the central nervous system (CNS), which facilitates neuronal conduction and general structural stabilization. HAPLN2 may also be involved in the formation of extracellular matrices, contributing to perineuronal nets and facilitating the understanding of a functional role of these extracellular matrices. HAPLN2 is found in several nuclei throughout the midbrain and hindbrain in a perineuronal net pattern.

## **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

## Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C for 12 months.  |
| Expiry Date:       | 12 months  |