antibodies .- online.com





anti-KERA antibody (AA 201-300) (AbBy Fluor® 488)



Go to Product page

()	۱ ۱	\cap	r	/1	\cap	۱ ۸	1
0	'V	ㄷ	ı١	νı	\Box	V	۷

Quantity:	100 μL
Target:	KERA
Binding Specificity:	AA 201-300
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KERA antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Keratocan
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	KERA
Alternative Name:	Keratocan (KERA Products)

Target Details

Bac	kar	ัดเ	ın	d	
Duo		\sim	<i>.</i>	u	•

Synonyms: CNA2, KERA, KERA_HUMAN, Keratan sulfate proteoglycan keratocan, Keratocan, KTN, SLRR2B.

Background: Small leucine-rich proteoglycans (SLRPs) such as Decorin, Biglycan, Fibromodulin, Keratocan, Lumican and Osteoglycin mediate extracellular matrix organization and are binding partners of TGF Beta. The Decorin core protein binds to growth factors, intercellular matrix molecules such as Fibronectin and Throm-bospondin, and to the Decorin endocytosis receptor. Fibromodulin is a collagen-binding keratan sulphate proteoglycan that influences adhesion processes of connective tissue and plays a role in fibrillogenesis by regulating collagen fibril spacing and thickness. Keratocan (KTN) develops corneal transparency and maintains the stromal matrix structure. Keratocan is a secreted protein in the extracellular matrix that binds to keratan sulfate chains. Keratocan is mainly detected in the cornea, but can also be expressed in trachea, intestine, ovary, lung and skeletal muscle. Defects in the gene encoding for Keratocan can cause cornea plana 2 (CNA2), an autosomal recessive disorder where the forward convex curvature of the cornea is flattened.

Pathways:

Glycosaminoglycan Metabolic Process

Application Details

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
\vdash	land	ling	
1 1	ıaııu	11111	۰

Expiry Date:

12 months