

[Go to Product page](#)

Datasheet for ABIN1389024

**anti-Netrin 3 antibody (AA 401-500) (Alexa Fluor 647)**

## Overview

Quantity:	100 µL
Target:	Netrin 3 (Ntn3)
Binding Specificity:	AA 401-500
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Netrin 3 antibody is conjugated to Alexa Fluor 647
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NTN3/Netrin-3
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Pig
Purification:	Purified by Protein A.

## Target Details

Target:	Netrin 3 (Ntn3)
Alternative Name:	NTN3/Netrin-3 ( <a href="#">Ntn3 Products</a> )

## Target Details

Background:	<p>Synonyms: NET3_HUMAN, Netrin-2-like protein, Netrin-3, NTN2L, NTN3.</p> <p>Background: Netrin proteins are a family of laminin-related secreted proteins that provide guidance signals for axonal growth and cell migration during development. Netrin signaling is dependent on the concentration of calcium outside the cell and the level of PKA activity. In axonal cells, a reduction in PKA activity converts the responsiveness of the axons to the netrin proteins as the cells are repelled, rather than attracted, by the netrin gradient. Neogenin serves as the primary guidance receptor for netrin-3. Netrin-2 and the corresponding mouse homolog netrin-3 are expressed primarily in the lower two-thirds of the spinal cord, and, like netrin-1, they can either attract or repel commissural axons at a distance. Netrin-3 proteins are associated with the axon fibers projecting from motor neurons and from neurons within sympathetic and sensory ganglia, suggesting that netrin-3 may be involved in pathfinding and fasciculation of axon projection. Neogenin serves as the primary guidance receptor for netrin-3. During peripheral nerve development, high netrin-3 expression has been detected in mesenchymal cells, sensory ganglia and muscles. In humans, the gene encoding for the netrin-3 protein is localized to chromosome 16p13.3.</p>
Pathways:	<a href="#">Regulation of Muscle Cell Differentiation</a>

## Application Details

Application Notes:	<p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p>
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

---

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months