# antibodies -online.com





## anti-Olfactomedin 3 antibody (AA 101-200) (FITC)



( )	1 /	$\sim$	KI /	110	Νę
	1//	$\vdash$	I \/	1 ←	٠// ٢

Quantity:	100 μL	
Target:	Olfactomedin 3 (OLFM3)	
Binding Specificity:	AA 101-200	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Olfactomedin 3 antibody is conjugated to FITC	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))	

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human OLFM3/Optimedin	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	Olfactomedin 3 (OLFM3)	
Alternative Name:	OLFM3/Optimedin (OLFM3 Products)	

#### **Target Details**

Racko	round:
Dacke	n Our iu.

Synonyms: Noe3, NOE3\_HUMAN, Noelin 3, Noelin-3, NOELIN3, NOELIN3 V1, NOELIN3 V2, NOELIN3 V3, NOELIN3 V4, NOELIN3 V5, NOELIN3 V6, Olfactomedin 3, Olfactomedin related ER localized protein 3, Olfactomedin-3, Olfactomedin3, Olfm3, Optimedin.

Background: OLFM3 is a 478 amino acid protein that interacts with myocilin. Myocilin is an extracellular protein that plays a key role in the actomyosin system and is responsible for controlling intraocular pressure. OLFM3 is a secreted protein that contains an olfactomedin-like (OLF) domain, an approximately 260 amino acid motif commonly found in secreted glycoproteins. OLFM3 localizes to the Golgi apparatus of the cell and is highly expressed in both eye and brain tissue. Mutations in the gene that encodes OLFM3 may cause severe glaucoma, a condition in which increased intraocular pressure within the eyeball causes loss of eye sight.

## **Application Details**

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

## Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	