antibodies -online.com





anti-FAT3 antibody (AA 601-800) (Biotin)



Overview

Quantity:	100 μL
Target:	FAT3
Binding Specificity:	AA 601-800
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAT3 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CDHF15/FAT3
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	FAT3
Alternative Name:	CDHF15/FAT3 (FAT3 Products)
Background:	Synonyms: Cadherin family member 15, CDHF15, CDHR10, FAT tumor suppressor homolog 3,

Fat3, FAT3_HUMAN, hFat3, Protocadherin Fat 3.

Background: The cadherins represent a family of Ca2+-dependent adhesion molecules that function to mediate cell to cell binding that is critical for the maintenance of structure and morphogenesis. Cadherins each contain a large extracellular domain at the N-terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. The relatively short C-terminal intracellular domain interacts with a variety of cytoplasmic proteins, including -catenin, to regulate cadherin function. The cadherin superfamily includes cadherins, protocadherins, desmogleins and desmocollins. FAT3 (FAT tumor suppressor homolog 3, also known as CDHF15 or CDHR10, is a 4,589 amino acid single-pass type I membrane protein expressed in ES cells, primitive neuroectoderm, fetal brain, infant brain, adult neural tissues and prostate. Containing thirty-three cadherin domains, four EGF-like domains and one laminin G-like domain, FAT3 may participate in the interactions between neurites derived from specific subsets of neurons during development.

Application Details

Application Notes:	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months