antibodies .- online.com





anti-Neurotrimin antibody (AA 101-200) (Alexa Fluor 350)



| () | 11/ | IN | /ie | A . |
|-----|-------------------|------|--------|--------|
| | / // | ۱ ات | / (− | ' \/\/ |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | Neurotrimin (NTM) | |
| Binding Specificity: | AA 101-200 | |
| Reactivity: | Human, Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This Neurotrimin antibody is conjugated to Alexa Fluor 350 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human Neurotrimin/HNT | |
|-----------------------|---|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse | |
| Predicted Reactivity: | Rat,Dog,Sheep,Pig,Horse,Rabbit | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | Neurotrimin (NTM) |
|-------------------|--------------------------------|
| Alternative Name: | Neurotrimin/HNT (NTM Products) |

Target Details

| Background: | Synonyms: hNT, IgLON family member 2, IGLON2, Neurotrimin, NT antibody NTM, | | | |
|---------------------|--|--|--|--|
| | NTRI_HUMAN. | | | |
| | Background: Cell adhesion molecules (CAMs) influence cell growth, differentiation, | | | |
| | embryogenesis, immune response and cancer metastasis by networking information from the | | | |
| | extracellular matrix to the cell. The four major families of cell adhesion molecules are | | | |
| | immunoglobulin (Ig) superfamily (calcium-independent transmembrane glycoproteins), | | | |
| | integrins (transmembrane non-covalently linked heterodimers of Alpha and Beta subunits), | | | |
| | calcium-dependent cadherins and divalent cation-dependent selectins. Regulation of neuronal | | | |
| | synaptic adhesion by CAMs has proven important for learning and memory. Proper embryonic | | | |
| | morphogenic development is also heavily dependent on the regulation of cell adhesion | | | |
| | molecules. Neurotrimin (hNT) is a neural cell adhesion molecule localizing to the cell | | | |
| | membrane, where it acts as a lipid-anchor. Neurotrimin belongs to the IgLON family of proteins, | | | |
| | a member of the larger immunoglobulin superfamily. | | | |
| Gene ID: | 50863 | | | |
| UniProt: | Q9P121 | | | |
| Application Details | | | | |
| Application Notes: | IF(IHC-P) 1:50-200 | | | |
| | IF(IHC-F) 1:50-200 | | | |
| | IF(ICC) 1:50-200 | | | |
| Restrictions: | For Research Use only | | | |
| Handling | | | | |
| Format: | Liquid | | | |
| Concentration: | 1 μg/μL | | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | | | |
| Preservative: | ProClin | | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | | | |
| Storage: | -20 °C | | | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | | | |
| | | | | |

| \vdash | land | lına |
|----------|-------|------|
| | iaria | шц |

Expiry Date:

12 months