antibodies -online.com





anti-Neurotrimin antibody (AA 101-200) (Biotin)



()	1 /	0	rv	/ 1 /	71	Α.
	1//	-	1 \/	16		1/1/
\sim	v	\sim	1 V	١,	_	v v

Quantity:	100 μL	
Target:	Neurotrimin (NTM)	
Binding Specificity:	AA 101-200	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Neurotrimin antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Neurotrimin/HNT	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Predicted Reactivity:	Rat,Dog,Sheep,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	Neurotrimin (NTM)
Alternative Name:	Neurotrimin/HNT (NTM Products)

Target Details

Background:	Synonyms: hNT, IgLON family member 2, IGLON2, Neurotrimin, NT antibody NTM,		
	NTRI_HUMAN.		
	Background: Cell adhesion molecules (CAMs) influence cell growth, differentiation,		
	embryogenesis, immune response and cancer metastasis by networking information from the		
	extracellular matrix to the cell. The four major families of cell adhesion molecules are		
	immunoglobulin (Ig) superfamily (calcium-independent transmembrane glycoproteins),		
	integrins (transmembrane non-covalently linked heterodimers of Alpha and Beta subunits),		
	calcium-dependent cadherins and divalent cation-dependent selectins. Regulation of neuronal		
	synaptic adhesion by CAMs has proven important for learning and memory. Proper embryonic		
	morphogenic development is also heavily dependent on the regulation of cell adhesion		
	molecules. Neurotrimin (hNT) is a neural cell adhesion molecule localizing to the cell		
	membrane, where it acts as a lipid-anchor. Neurotrimin belongs to the IgLON family of proteins,		
	a member of the larger immunoglobulin superfamily.		
Gene ID:	50863		
UniProt:	Q9P121		
Application Details			
Application Notes:	WB 1:300-5000		
	IHC-P 1:200-400		
	IHC-F 1:100-500		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C for 12 months.		

	_	
-	II	1:
-	-222	una
-	Hand	1111()

Expiry Date:

12 months