# antibodies -online.com





# anti-Junctophilin 3 antibody (AA 181-280) (Alexa Fluor 488)



Go to Product page

| $\sim$ |     |     |     |
|--------|-----|-----|-----|
|        | N/P | r\/ | i⊢₩ |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | Junctophilin 3 (JPH3)  |
| Binding Specificity: | AA 181-280   |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This Junctophilin 3 antibody is conjugated to Alexa Fluor 488  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human Junctophilin 3 |
|-----------------------|--|
| Isotype:              | IgG  |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Zebrafish          |
| Purification:         | Purified by Protein A.   |

## **Target Details**

| Target:   | Junctophilin 3 (JPH3)          |
|---|--------------------------------|
| Alternative Name:   | Junctophilin 3 (JPH3 Products) |
| Background: Synonyms: JP-3, JP3, JPH3, JPH3_HUMAN, Junctophilin3, Junctophilin type 3, Junctophilin |                                |

TNRC22, Trinucleotide repeat-containing gene 22 protein.

Background: Junctophilins are components of the junctional complexes between plasma membranes and endoplasmic or sarcoplasmic reticulums present in all excitable cells. Junctophilins contain a cytoplasmic domain which binds to the plasma membrane, as well as an ER/SR membrane spanning hydrophobic C-terminal segment. The three subtypes in this family are Junctophilin-1, -2 and -3. Junctophilin-1 is predominantly expressed in skeletal muscle, but is also expressed at low levels in heart. Junctophilin-2 is expressed in heart and skeletal muscle. Mutant mice lacking the Jph2 gene exhibit embryonic lethality and possess cardiac myocytes that express abnormal calcium transients. Junctophilin-3 is expressed in brain. The JPH3 alternatively spliced exon 2A has been suggested as a site for CTG repeat expansion leading to a Huntington disease-like autosomal dominant disorder.

## **Application Details**

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--------------------|
|                    | IF(IHC-F) 1:50-200 |
|                    | IF(ICC) 1:50-200   |

Restrictions: For Research Use only

#### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |