# antibodies -online.com





# anti-Junctophilin 3 antibody (AA 181-280) (Biotin)



Go to Product page

( )	11/0	K\ /	iew	1
	$\cup$	ועוי	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	Junctophilin 3 (JPH3)
Binding Specificity:	AA 181-280
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Junctophilin 3 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Junctophilin 3
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Chicken, Zebrafish
Purification:	Purified by Protein A.

## **Target Details**

Torget	Junatankilia 2 / IDU2\	
Target:	Junctophilin 3 (JPH3)	
Alternative Name:	Junctophilin 3 (JPH3 Products)	
Background:	und: Synonyms: JP-3, JP3, JPH3, JPH3, HUMAN, Junctophilin3, Junctophilin type 3, Junctophilin	

TNRC22, Trinucleotide repeat-containing gene 22 protein.

Background: Junctophilins are components of the junctional complexes between plasma membranes and endoplasmic or sarcoplasmic reticulums present in all excitable cells. Junctophilins contain a cytoplasmic domain which binds to the plasma membrane, as well as an ER/SR membrane spanning hydrophobic C-terminal segment. The three subtypes in this family are Junctophilin-1, -2 and -3. Junctophilin-1 is predominantly expressed in skeletal muscle, but is also expressed at low levels in heart. Junctophilin-2 is expressed in heart and skeletal muscle. Mutant mice lacking the Jph2 gene exhibit embryonic lethality and possess cardiac myocytes that express abnormal calcium transients. Junctophilin-3 is expressed in brain. The JPH3 alternatively spliced exon 2A has been suggested as a site for CTG repeat expansion leading to a Huntington disease-like autosomal dominant disorder.

## **Application Details**

Application Notes:	WB 1:300-5000
application (40 teo.	VVD 1.000 0000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months