antibodies .- online.com







anti-LSAMP antibody (AA 51-150) (FITC)



()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Quantity:	100 μL	
Target:	LSAMP	
Binding Specificity:	AA 51-150	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This LSAMP antibody is conjugated to FITC	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human LSAMP	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	LSAMP	
Alternative Name:	LSAMP (LSAMP Products)	
Background:	ground: Synonyms: IgLON family member 3, IGLON3, Lam, LAMP, Limbic system associated men	

protein, Limbic system associated membrane protein precursor, Limbic system-associated membrane protein, LSAMP, LSAMP_HUMAN.

Background: LSAMP is a neuronal surface glycoprotein found in cortical and subcortical regions of the limbic system expressed on the surface of somata and proximal dendrites of neurons. It is a member of the immunoglobulin (Ig) superfamily and belongs to the IgLON subfamily of cell adhesion molecules. LSAMP contains three Ig domains and a glycosylphosphatidylinositol anchor. It is a highly conserved protein between rodents and humans (99 % sequence identity) and it is involved in the regulation of neurite outgrowth and mediation of proper circuit formation of limbic pathways. This suggests that LSAMP plays an important role in the development and function of the limbic system. In addition, LSAMP may function as a tumor suppressor in renal carcinomas.

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	