antibodies -online.com





anti-PCDH11X antibody (AA 601-700) (Biotin)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	PCDH11X (PCDH11)	
Binding Specificity:	AA 601-700	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PCDH11X antibody is conjugated to Biotin	
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PCDH11X
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	PCDH11X (PCDH11)
Alternative Name: PCDH11X (PCDH11 Products)	
Background: Synonyms: PCDH11, PCDH22, PCDHX, PCDHY, Protocadherin 11, Protocadherin 11 X Ii	

Protocadherin 11 Y linked, Protocadherin 22, Protocadherin on the X chromosome, Protocadherin on the Y chromosome, Protocadherin PC, Protocadherin prostate cancer, PC11X_HUMAN.

Background: This gene belongs to the protocadherin gene family, a subfamily of the cadherin superfamily. The encoded protein consists of an extracellular domain containing 7 cadherin repeats, a transmembrane domain and a cytoplasmic tail that differs from those of the classical cadherins. The gene is located in a major X/Y block of homology and its Y homolog, despite divergence leading to coding region changes, is the most closely related cadherin family member. The protein is thought to play a fundamental role in cell-cell recognition essential for the segmental development and function of the central nervous system.

Transcripts arising from alternative splicing encode isoforms with variable cytoplasmic domains. [provided by RefSeq, Jul 2008].

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months