antibodies -online.com





anti-Protocadherin gamma Subfamily C, 3 (PCDHGC3) (AA 251-350) antibody (Biotin)



()	1 /	0	rv	/ 1 /	71	Α.
	1//	\vdash	1 \/	16		1/1/
\sim	v	\sim	1 V	١,	_	v v

Quantity:	100 μL	
Target:	Protocadherin gamma Subfamily C, 3 (PCDHGC3)	
Binding Specificity:	AA 251-350	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PCDHGC3/PCDH2		
Isotype:	IgG		
Predicted Reactivity:	Human,Mouse,Rat,Horse		
Purification:	Purified by Protein A.		

Target Details

Target: Protocadherin gamma Subfamily C, 3 (PCDHGC3)		
Alternative Name:	PCDHGC3/PCDH2 (PCDHGC3 Products)	
Background: Synonyms: Cadherin like 2. PC 43. PC43. PCDH 2. PCDH gamma C3. PCDH2. PC		

Protocadherin 2, Protocadherin 43, Protocadherin gamma C3, Protocadherin gamma subfamily C 3, Protocadherin43, PCDGK_HUMAN.

Background: Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin gene clusters, designated alpha, beta and gamma, all of which contain multiple tandemly arranged genes. PCDH2 (protocadherin-2), also known as PCDHGC3 (protocadherin gamma subfamily C, 3) or PC43, is a 934 amino acid single-pass type I membrane protein that contains six cadherin domains and belongs to the protocadherin gamma family. Functioning as a calcium-dependent cell-adhesion protein, PCDH2 is thought to be involved in the establishment and maintenance of neuronal connections within the brain. Multiple isoforms of PCDH2 exist due to alternative splicing events.

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months